Detection of a back-scattering region at 108 km height during sunset

C. M. Denardini, M. A. Abdu, E. R. de Paula, C. M. Wrasse, J. H. A. Sobral Instituto Nacional de Pesquisas Espaciais - P.O. 515 - S. J. Campos, SP, Brazil

Abstract


We have obtained Range Time Intensity (RTI) maps covering the equatorial electrojet heights during daytime and evening in the Brazilian sector using the RESCO 50 MHz back-scatter radar (2.33° S, 44.2° W, DIP: -0.5), at São Luís. These maps revealed the appearance of a scattering region at about 108 km of height during the sunset period. It was observed mainly during summer period under quiet magnetic activities conditions. Back-scatter coherent radars are sensitive to field aligned meter-scale plasma irregularities and the type of 3-m irregularity region we have observed, and we present here has not been reported before to our best of knowledge. The occurrence of this echo region coincides in local time with the maximum intensity of evening pre reversal eastward electric field of the F region ionosphere. Theory of the divergence of the equatorial electrojet current in the evening jonosphere has been proposed (Haerendel and Eccles, 1992, JGR) to explain its partial contribution to the development of the pre reversal electric field. The theory predicts enhanced zonal electric field and hence vertical electric field below 300 km as necessary as consequence of the electrojet divergence in the evening. In this paper the experimental results of the enhanced echoes from higher heights of the EEJ region do seem to provide an evidences that the divergence of the equatorial electrojet current can indeed be the driver of the scattering region observed. For the present study we have selected some days of radar observation of the E region heights during summer under quiet and disturbed magnetic condition. We also have obtained the amplitude of F-region plasma drift (pre reversal enhancement) over the same observatory, deduced from hmF2 obtained by the digisonde.

Acknowledgements

This work was supported by FAPESP (Fundação de Amparo a Pesquisa de Estado de São Paulo) through thematic project grant no. 99/00437. Support received through CNPq (Conselho Nacional de Pesquisa e Desenvolvimento) grants nos. 520185/95-1 and 522919/96-0 is also acknowledged. C. M. D. wishes to thank FAPESP for the financial support to his doctoral degree program through the grant n° 98/16156-8, and for the financial support to his post-doctoral program through the grant n° 03/01146-7.

		IAGA 2005
Programme Sections	14:40 - 15:00	IAGA2005-A-00045 Shirochkov, A.V.; Makarova, L.N.; Sokolov, S.N.; Sheldon, W.R. lonospheric Effects of the Global Geomagnetic Storms
▶ IAGA & ICMA Joint Symposia	15:00	COFFEE BREAK
► ICMA Symposium		
▶ Division I	Chairperson:	N.N.
▶ Division II		
▶ Division III		<u>IAGA2005-A-00971</u> Basu, S. ; Basu, Su.; Groves, K.; Mackenzie, E.; Sheehan, R.; Rich, F.; Keskinen, M. Plasma structuring in the ionosphere at middle and equatorial latitudes during magneti superstorms
▶ Division IV		
▶ Division ∨		
Inter-Divisional Commission on Developing Countries Inter-Divisional Commission on History Inter-Divisional Inter-Divisiona	15:40 - 16:00	IAGA2005-A-00062 Denardini, C. M.; Abdu, M. A.; de Paula, E. R.; Wrasse, C. M.; Sobral, J. H. Detection of a back-scattering region at 108 km height during sunset
		AGA2005-A-00845 Veenadhari, B; Alex, S Ionospheric response to intense magnetic storms of 29 October and 20 November, 2003.
	16:20 - 16:40	IAGA2005-A-00214 Liu, RY.; Xu, ZH.; Liu, SL.; Zhang, BC.; Hu, HQ.; Qi, X. Ionospheric variations in the F2-layer at Zhongshan Station, Antarctica
		IAGA2005-A-00639 Hady, A.; Shaltout, M. Historical data of geomagnetic storms and climatic change
	17:00	END OF SESSION





10" Scientific Assembly of the International Association of Geomagnetism and Aeronomy Toulouse - France, July 18-29, 2005