



The solar origin of the October 21st-22nd (1999) very intense geomagnetic storm

A. Dal Lago, L. E. A. Vieira, E. Echer, F. L. Guarnieri, M. R. da Silva, A. de Lucas, W. D. Gonzalez, A. L. Clua de Gonzalez

Instituto Nacional de Pesquisas Espaciais, INPE, São José dos Campos, SP, Brazil.

Copyright 2005, SBGf - Sociedade Brasileira de Geofísica

This paper was prepared for presentation at the 9th International Congress of the Brazilian Geophysical Society held in Salvador, Brazil, 11-14 September 2005.

Contents of this paper were reviewed by the Technical Committee of the 9th International Congress of the Brazilian Geophysical Society. Ideas and concepts of the text are authors' responsibility and do not necessarily represent any position of the SBGf, its officers or members. Electronic reproduction or storage of any part of this paper for commercial purposes without the written consent of the Brazilian Geophysical Society is prohibited.

Abstract

In this work we address the solar origin of the October 21st-22nd (1999) very intense geomagnetic storm, which was caused by the interaction between an interplanetary ejecta (ICME) and a high speed stream. We believe that a high speed stream compressed the interplanetary ejecta and intensified its internal magnetic field, thus increasing the intensity of the geomagnetic storms. This kind of event is very difficult to predict based on remote observations of the solar corona, such as those from the Large Angle and Spectroscopic Coronagraph – LASCO, aboard the Solar and Heliospheric Observatory – SOHO, because they are normally related to relatively slow CMEs. Using combined data from different instruments, we present evidences on the solar disk of a coronal hole just beside (to the east) the active region from which the coronal mass ejection was lifted off. In the last solar cycle at least 1 out of 14 very intense geomagnetic storms, i.e. peak Dst < -200 nT, was caused by such a mechanism.

Introduction

High speed streams, originated in coronal holes, are often observed following interplanetary coronal mass ejection – ICMEs, at 1 AU (Klein and Burlaga, 1982). One of the most important effects of the interaction between these two structures is the intensification of the magnetic cloud field in the latter part, due to compression, thus leading to a stronger geomagnetic perturbation if the cloud's intensified magnetic field is pointing southward (Burlaga et al., 1987; Bothmer and Schwenn, 1995; Badruddin, 1998; Fenrich and Luhmann, 1998; Dal Lago et al., 2001, 2002).

One of the aims of space weather studies is to forecast earth disturbances using solar observation in order to have enough time to prevent damages in satellites and other man made equipment. It is well known that coronal mass ejections – CMEs, observed by coronagraphs at the sun are one of the most important causes of geomagnetic disturbances (e. g. Gosling, 1993).

In this paper we present the solar origin of the October 21st-22nd (1999) very intense geomagnetic storm, which

was caused by the interaction between an interplanetary ejecta (ICME) and a high speed stream. Specifically, we address the solar causes of this event, using combined data from different instruments, both space and ground based.

Method

Since 1996, the Large Angle and Spectroscopic Coronagraph – LASCO (Brueckner et al., 1995), and the Extreme ultraviolet Imaging Telescope – EIT, both aboard the Solar and Heliospheric Observatory – SOHO provide continuous measurements of CMEs and their disk counterparts, like EIT eruptions. LASCO is a white light coronagraph and it is able to reveal eruptions of coronal mass ejections. One of the EIT filters is able to image the solar disk in 19.5 nm with enough time resolution to identify EIT eruptions related to CMEs. The National Solar Observatory – NSO, at Kitt Peak, provides coronal hole maps, obtained by ground based instrumentation, which detect the 1083 He I absorption line (Harvey et al., 1975). To complete this set of solar observations, we have in situ magnetic field and plasma data from the Advanced Composition Explorer – ACE satellite, siting in the Lagrangean point L1, and Kyoto Dst index, which show the arrival at earth of a complex interplanetary structure and a very intense geomagnetic storm, respectively.

Solar origin of the October 21-22 (1999) disturbance

On the October 18th, 1999, at 00:06 UT, LASCO C2 observed a coronal mass ejection, which evolved to a halo in LASCO C3 instrument with an expansion speed of 546.1 km/s, as shown in Figure 1. Associated with this CME, an EIT eruption was observed toward the east side of the solar disk, during the last hour of day 17. Its evolution can be seen in Figure 2.

NSO/KP maps from October 17th to October 18th, as well as YOHKOH Soft X-ray telescope on October 19th, clearly reveal a positive coronal hole close to the site of the EIT eruption observed in the EIT images of day 17, as shown in Figure 3.




Fig. 1 – LASCO C2 (top images) and LASCO C3 (bottom) observations of the October 18th (1999) CME.




Fig. 2 – EIT running difference images of the eruption during the last hour of day 17, in October, 1999. Another eruption in the southern solar hemisphere is also visible in the image, but this one was not related with our event.





Fig. 3 – NSO/KP coronal hole maps (top images) from Oct. 17th and Oct. 18th, 1999. Yohkoh SXT image from Oct. 19th. In all images, a clear (positive) coronal hole can be identified.

Combining the observation from EIT in Figure 2 from Oct. 17th, at 23:24UT with the NOS/KP coronal hole map of Oct. 17th, one can easily see that the coronal hole was sitting beside the EIT eruption site, more precisely to the east, as shown in Figure 4.




Fig. 4 – Combined observations from SOHO/EIT 195 and NOS/KP coronal hole map, from Oct. 17th, 1999.

Close to the earth, at L1, an interplanetary shock was detected by the ACE magnetic field and plasma instruments, 73.5 hours after the first appearance of the related CME in the LASCO C2 field-of-view, on October 21st (1999), at 01:34 UT, as shown in Figure 5.




Fig. 5 – From top to bottom: Magnetic field intensity B and its Bz component, By and Bx components, solar wind velocity and density, and the Dst index.

Immediately after the ICME structure (second vertical dotted line), a high speed stream can be seen, with a speed close to 700 km/s. The interaction between this high speed stream and the ICME probably caused compression and intensification of the magnetic field inside the ICME, as can be observed in the top panel of Figure 5. The very intense negative Bz field in the rear portion of the ICME structure lead to the very intense geomagnetic storm.

Summary and Conclusions

We have presented the solar origin of the October 21st-22nd (1999) very intense geomagnetic storm, which was caused by the interaction between an interplanetary ejecta (ICME) and a high speed stream. Combined observations from SOHO/EIT 195 and NSO/KP maps revealed a proximity of an eastward positive coronal hole and an EIT eruption, both observed on the Oct. 17th. Following that, on the Oct. 18th SOHO/LASCO observed a CME, closely related in time with the EIT eruption. The combination of these solar structures was probably the solar origin of the ICME-high speed stream structure

observed by ACE instruments near earth, on the Oct. 21st-22nd, and hence the cause of the very intense geomagnetic storm.

Acknowledgments

The authors would like to acknowledge: LASCO and EIT workteams for providing the coronagraph and ultraviolet observations used in this work; the National Solar Observatory/Kitt Peak, under the National Science Foundation support, for providing the coronal hole maps; the YOHKOH SXT workteam for the soft x-ray image shown in this work; N. Ness (Bartol Research Institute), D. J. McComas (Southwest Research Institute), R. Lepping (NASA/GSFC), K. Ogilvie (NASA GSFC) and CDAWeb for interplanetary magnetic field and plasma data; and Kyoto WDC for the Dst index. SOHO is an international cooperation between ESA and NASA. The authors would also like to acknowledge Fundação de Amparo a Pesquisa do Estado de São Paulo – FAPESP of Brazil, for supporting partially this work under the project 02/12723-2, and Conselho Nacional de Desenvolvimento Científico e Tecnológico – CNPq, of Brazil, for supporting partially this work.

References

- Badruddin, Interplanetary Shocks, Magnetic Clouds, Stream Interfaces and Resulting Geomagnetic Disturbances, *Planetary Space Science*, 46, 1015-1028, 1998.
- Bothmer, V., and R. Schwenn, The Interplanetary and Solar Causes of Major Geomagnetic Storms, *J. Geomag. Geoelectr.*, 47, 1127-1132, 1995.
- Brueckner, G. E.; Howard, R. A.; Koomen, M. J.; Korendyke, C. M.; Michels, D. J.; Moses, J. D.; Socker, D. G.; Dere, K. P.; Lamy, P. L.; Llebaria, A.; Bout, M. V.; Simnett, G. M.; Bedford, D. K.; Eyles, C. J. The large angle spectroscopic coronagraph (LASCO). *Solar Phys.* v.162, p.357-402, 1995.
- Burlaga, L. F., Magnetic Clouds and Force-Free with Constant Alpha, *J. Geophys. Res.*, 93, 7217-7224, 1988.
- Burlaga, L. F., K. W. Behannon, and L. W. Klein, Compound Streams, Magnetic Clouds, and Major Geomagnetic Storms, *J. Geophys. Res.*, 92, 5725-5734, 1987.
- Dal Lago, A., W. D. Gonzalez, A. L. Clua de Gonzalez, and L. E. A. Vieira, Compression of Magnetic Clouds in Interplanetary Space and Increase in their Geoeffectiveness, *J. Atm. Solar-Terr. Phys.*, 63, 451-455, 2001.
- Dal Lago, A., W.D. Gonzalez, A.L.C. de Gonzalez, e L.E. Vieira, Stream-interacting magnetic clouds causing very intense geomagnetic storms. *Adv. Space Res.*, v.30, n.10, p.2225-2229, 2002.

Fenrich, F. R., and J. G. Luhmann, Geomagnetic Response to Magnetic Clouds of Different Polarity, *J. Geophys. Res.*, 25, 2999-3002, 1998.

Harvey, J. W.; Krieger, A. S.; Tomothy, A. F.; Vaiana, G. S.; *Bull American Astron. Soc.*, 7, 358, 1975.

Klein, L. W., and L. F. Burlaga, Interplanetary Magnetic Clouds at 1 AU, *J. Geophys. Res.*, 87, 613-624, 1982.

Gosling, J. T, The Solar Flare Myth, *J. Geophys. Res.*, 98, 18937-18949, 1993.